

EXPRESS MAIL NUMBER
EL92348381015

**RAPIDLY DEGRADING GFP-FUSION PROTEINS
AND METHODS OF USE**

5

BACKGROUND OF THE INVENTION

Cross-Reference to Related Application

10
15 This application is a continuation in part of U.S. patent
application Serial Number 09/062,102, filed April 17, 1998.

Field of the Invention

20 This invention relates to the field of biochemical assays
and reagents. More specifically, this invention relates to modified
fluorescent proteins and to methods for their use.

Description of the Related Art

Because of its easily detectable green fluorescence, green fluorescent protein (GFP) from the jellyfish *Aequorea victoria* has been used widely to study gene expression and protein localization. GFP fluorescence does not require a substrate or cofactor; hence, it is possible to use this reporter in numerous species and in a wide variety of cells. GFP is a very stable protein which can accumulate and thus is often toxic to mammalian cells.

Recently, crystallographic structures of wild-type GFP and the mutant GFP S65T reveal that the GFP tertiary structure resembles a barrel (Ormo et al. (1996) *Science* 273: 1392-1395; Yang, F., Moss, L. G., and Phillips, G. N., Jr. (1996) *Nature Biotech* 14: 1246-1251). The barrel consists of beta sheets in a compact antiparallel structure. In the center of the barrel, an alpha helix containing the chromophore is shielded by the barrel. The compact structure makes GFP very stable under diverse and/or harsh conditions, such as protease treatment, making GFP an extremely useful reporter in general. On the other hand, its stability makes it difficult to determine short-term or repetitive events.

A great deal of research is being performed to improve the properties of GFP and to produce GFP reagents useful for a

variety of research purposes. New versions of GFP have been developed via mutation, including a "humanized" GFP DNA, the protein product of which has increased synthesis in mammalian cells (see Cormack, et al., (1996) *Gene* 173, 33-38; Haas, et al., (1996) *Current Biology* 6, 315-324; and Yang, et al., (1996) *Nucleic Acids Research* 24, 4592-4593). One such humanized protein is "enhanced green fluorescent protein" (EGFP). Other mutations to GFP have resulted in blue-, cyan- and yellow-fluorescent light emitting versions.

Ornithine decarboxylase (ODC) is an enzyme critical in the biosynthesis of polyamines. Murine ornithine decarboxylase is one of most short-lived proteins in mammalian cells, with a half life of about 30 minutes (Ghoda, et al., (1989) *Science* 243, 1493-1495; and Ghoda, et al. (1992) *Mol. Cell. Biol.* 12, 2178-2185). Rapid degradation of murine ornithine decarboxylase has been determined to be due to the unique composition of its C-terminus, a portion of which has a PEST sequence—a sequence which has been proposed as characterizing short-lived proteins. The PEST sequence contains a region enriched with proline (P), glutamic acid (E), serine (S), and threonine (T), often flanked by basic amino acids, lysine, arginine, or

histidine (Rogers, et al., (1989) *Science* 234:364-68; Reichsteiner, M. (1990) *Seminars in Cell Biology* 1:433-40).

The ornithine decarboxylase of *Trypanosoma brucei* (TbODC) does not have a PEST sequence, and is long-lived and quite
5 stable when it is expressed in mammalian cells (Ghoda, et al.. (1990) *J. Biol. Chem.* 265: 11823-11826); however appending the Cterminus of murine ornithine decarboxylase to TbODC makes TbODC become unstable. Moreover, deletion of the C-terminal, PEST-containing region from murine ornithine decarboxylase prevents its rapid
10 degradation (Ghoda, L., et al. (1989) *Science* 243: 1493-1495).

The prior art is deficient in a destabilized or short-lived GFP. The present invention fulfills this need in the art.

SUMMARY OF THE INVENTION

A rapid turnover or destabilized GFP can be used in research applications where prior art GFPs cannot. Such applications include using the destabilized GFP as a genetic reporter for analyzing
20 transcriptional regulation and/or cis-acting regulatory elements, or as a tool for studying protein degradation. Further, a rapid turnover

GFP permits easier development of stable cell lines which express the GFP gene, since toxic levels of GFP are avoided because the GFP protein is degraded quickly.

The present invention provides a fusion protein with a half life decreased markedly from that of wildtype GFP. In one embodiment, there is provided a fusion protein comprising an EGFP fused to a peptide producing a destabilized protein. In another embodiment, there is provided a fusion protein with a half life of about ten hours or less, preferably with a half life of about 4 hours or less, more preferably with a half life of 2 hours or less and even more preferably with a half life of 1 hour or less. A preferred embodiment of this aspect of the invention includes EGFP, and/or a PEST sequence-containing portion of a C-terminus of murine ornithine decarboxylase (MODC). Specific preferred embodiments of the present invention include EGFP-MODC₃₇₆₋₄₆₁; EGFP-MODC₃₇₆₋₄₅₆; EGFP-MODC₄₂₂₋₄₆₁; P426A/P427A; P438A; E428A/E430A/E431A; E444A; S440A; S445A; T436A; D433A/D434A; and D448A.

In yet another aspect of the invention, there is provided an isolated DNA molecule encoding a fluorescent fusion protein with a half life that is markedly decreased from that of wildtype GFP. In

one embodiment of this aspect of the invention, there is provided an isolated DNA molecule encoding a fluorescent fusion protein with a half life of about ten hours or less, preferably with a half life of about 4 hours or less, more preferably with a half life of 2 hours or less and even more preferably with a half life of 1 hour or less. In a preferred embodiment of this aspect of the invention, the isolated DNA molecule encoding the fluorescent fusion protein is a synthetic GFP gene containing codons preferentially found in highly expressed human proteins. Further, the present invention provides a vector capable of expressing the isolated DNA molecule encoding a GFP fusion protein with a decreased half life. In one embodiment of the vector, the vector contains an inducible promoter.

In another aspect of the invention, there is provided a method of labeling cells with a transient GFP reporter. In this method, a DNA vector comprising an inducible promoter and the isolated DNA encoding a GFP fusion protein with a decreased half life is utilized. This vector is transfected into cells which are cultured under conditions in which the promoter induces transient expression of the GFP fusion protein of the present invention, which provides a transient fluorescent label.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic map of EGFP and EGFP-MODC fusion proteins. EGFP is fused with a region of the C terminus of MODC, including amino acids from 376 to 461, 376 to 456 or 422 to 461. The fusion proteins were expressed in CHO K1 Tet-off cells and their fluorescence intensities were compared under a fluorescence microscope.

Figure 2 shows the fluorescence stabilities of EGFP and EGFP-MODC₄₂₂₋₄₆₁ in cells in the presence of cycloheximide and examined with a fluorescence microscope. CHO K1 Tet-off cells were transfected with vectors expressing these two proteins. After 24 hours, the transfected cells were treated with 100 mg/ml cycloheximide for 0, 1, 2, 3, and 4 hours.

Figure 3 shows flow cytometric analysis of the fluorescence stabilities of EGFP and EGFP-MODC₄₂₂₋₄₆₁. CHO K1 Tet-off cells were transfected with EGFP and EGFP-MODC₄₂₂₋₄₆₁. After 24 hours, the transfected cells were treated with 100 µg/ml cycloheximide for 0, 1, 2, and 3 hours. The treated cells were collected with EDTA and 10,000 cells were subjected to FACS analysis

Figure 4 is a graph summarizing the flow cytometric data from figure 3, demonstrating that EGFP-MODC₄₂₂₋₄₆₁-transfected cells rapidly lose fluorescence after cycloheximide treatment, whereas EGFP cells maintain fluorescence.

5 **Figure 5** is a photograph of western blot analysis of protein stabilities of EGFP and EGFP-MODC₄₂₂₋₄₆₁. Cells collected during flow cytometry were used for preparing cell lysates. The cell lysates were subject to SDS gel electrophoresis and transferred onto a membrane. EGFP and the EGFP fusion protein were detected with a
10 monoclonal antibody against GFP.

Figure 6 is a schematic map of the PEST sequence of the fusion EGFP-MODC₄₂₂₋₄₆₁ indicating the position of the mutations.

Figure 7 is a table summarizing the results obtained measuring persistence of fluorescent signal in transfected CHO K1
15 Tet-off cells expressing EGFP, EGFP-MODC₄₂₂₋₄₆₁, and the PEST mutants. Transfection was performed in CHO/tTA cells using the procedure given in Example 2. After 24 hours, cells were treated with cycloheximide for 0, 2, and 4 hours, and analyzed for fluorescence by FACS Caliber.

Figure 8 shows a schematic illustration of d2EGFP, dECFP and dEYFP.

Figure 9 shows the construction of destabilized EGFP Variants.

5 Figure 10 shows the fluorescence stabilities of EGFP and dEGFP Variants.

Figure 11 shows the increase in induction by CRE-d1EGFP and CRE-d2EGFP.

10 DETAILED DESCRIPTION OF THE INVENTION

The invention describes a genetically-engineered fluorescent protein that is destabilized, having a rapid turnover in a cell. More specifically, this fusion protein comprises a fluorescent protein which has a half life of no more than about ten hours and most preferably with a half life of no more than 2 hours. Preferably, the fluorescent protein is selected from the group consisting of EGFP, ECFP and EYFP. In one embodiment, the engineered GFP is a fusion protein of EGFP and a peptide the inclusion of which produces a destabilized protein. An example of such a peptide is the C-terminal region of murine

15

20

ornithine decarboxylase (MODC). In a specific, illustrative case, the degradation domain of murine ornithine decarboxylase from amino acids 422 to 461 was appended to the C-terminal end of an enhanced variant of GFP (EGFP). The fluorescence intensity of the EGFP-MODC₄₂₂₋₄₆₁ fusion protein in transfected cells was similar to that of EGFP, but the fusion protein, unlike EGFP, dissipated over time in cells treated with cycloheximide. The half-life of the fluorescence of the EGFP-MODC₄₂₂₋₄₆₁ fusion protein was about 2 hours, while that of EGFP was more than 24 hours. The ornithine decarboxylase degradation domain dramatically decreases EGFP stability.

The rapid turnover version of EGFP has at least four advantages over EGFP. The rapid turnover of the EGFP-MODC fusion causes less toxicity to cells expressing the fusion protein. Thus, one advantage is the feasibility of establishing stable cell lines using DNA coding for EGFP-murine ornithine decarboxylase. Further, the destabilized EGFP-MODC decreases EGFP accumulation. Accumulation of fluorescent protein can interfere with the sensitivity of analysis. Thus, the destabilized, rapid turnover fusion protein renders more sensitive results. Additionally, destabilized EGFP can be used as a transient reporter to study transcriptional regulation and/or action of cis-acting regulatory elements. Finally, the EGFP-MODC fusion

protein can be used to study processes involving multiple gene expression.

Moreover, the EGFP-MODC fusion protein has the advantages inherent to use of EGFP. For example, the use of EGFP in drug screening assays is particularly advantageous because GFP fluorescence can be detected intracellularly without performing additional expensive steps; e.g. lysing cells, adding exogenous substrates or cofactors, fixing the cell preparation, etc. A single illustration of such an assay is screening test compounds for interruption of the TNF activation pathway, a pathway which ultimately affects apoptosis. Compounds identified in the assay would be useful in controlling the cellular processes involved in cancer and inflammation.

Further, the reporter gene of the present invention can be linked with different enhancer elements and used to monitor diverse biological processes such as heat response, response to heavy metals, glucocorticoid activation or response to cAMP. In particular, destabilized EGFP is useful for studying developmental processes where genes are transiently expressed, dynamics of protein transport, localization of proteins within cells, and periodic and cyclical expression of genes that control unique biological phenomena

such as circadian rhythms. Indeed, other applications of the EGFP-MODC fusion protein in screening assays would be appreciated readily by those having ordinary skill in this art.

Moreover, by using an inducible promoter, expression of the EGFP-MODC fusion protein is activated or deactivated at will, making a construct expressing the protein useful in cell lineage studies. Prior art GFP models express GFP at levels that are toxic and interfere with cell development, thus making cell lineage studies impossible. Additionally, destabilized EGFP can be used as a reporter to study the kinetics of mRNA transcription from a regulated promoter, because the fluorescence intensity of destabilized EGFP is a direct measure of the level of gene expression at any given time point.

As used herein, the term "GFP" refers to the basic green fluorescent protein from *Aequorea victoria*, including prior art versions of GFP engineered to provide greater fluorescence or fluoresce in different colors. The sequence of *A. victoria* GFP has been disclosed in Prasher D.C. et al. (1992) *Gene* 111:229-33.

As used herein, the term "EGFP" refers to GFP which has been "humanized", as reported in Kain et al. (1995) *Biotechniques* 19(4):650-55. "Humanized" refers to changes made to the GFP

nucleic acid sequence to optimize the codons for expression of the protein in human cells.

As used herein, the term "peptide which produces a destabilized protein" refers to a sequence of amino acids or a peptide which promotes destabilization or rapid turnover of the protein of which it is a part; i.e., by inducing protein degradation. The PEST sequence described herein is one such sequence. Other sequences known in the art are those peptides that promote phosphorylation and protein-protein interactions.

As used herein, the term "EGFP-MODC" refers to EGFP fused at its C terminus to murine ornithine decarboxylase sequences.

As used herein, the term "P438A" refers to an EGFP-MODC fusion protein in which the proline at position 438 in the murine ornithine decarboxylase sequence (a proline residing in the PEST portion of the sequence) has been replaced with alanine. The same nomenclature is used for EGFP-MODC mutants P426A/P427A; E428A/E430A/E431A; E444A; S440A; S445A; T436A; D433A/D434A; and D448A. Further elucidation is shown in Figure 6.

As used herein, the term "half life" refers to the period of time in which half of the fluorescent signal from a fluorescent protein expressed in cells disappears and half remains.

As used herein, the term "Tc" refers to tetracycline.

As used herein, the term "CHX" refers to cycloheximide.

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Maniatis, Fritsch & Sambrook, "Molecular Cloning: A Laboratory Manual (1982); "DNA Cloning: A Practical Approach," Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" (B.D. Hames & S.J. Higgins eds. (1985)); "Transcription and Translation" (B.D. Hames & S.J. Higgins eds. (1984)); "Animal Cell Culture" (R.I. Freshney, ed. (1986)); "Immobilized Cells and Enzymes" (IRL Press, (1986)); B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

A "DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in either single stranded form or a double-stranded helix. This term refers only to the primary and secondary structure of the molecule,

and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, *inter alia*, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes.

5 A DNA "coding sequence" is a DNA sequence which is transcribed and translated into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (e.g., mammalian) DNA, and synthetic DNA sequences. A polyadenylation signal and transcription termination sequence may be located 3' to the coding sequence.

15 Transcriptional and translational control sequences are DNA regulatory sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide for and/or regulate expression of a coding sequence in a host cell.

20 A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For

purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels
5 detectable above background. Within the promoter sequence will be found a transcription initiation site, as well as protein binding domains responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes. Various promoters, including inducible promoters, may
10 be used to drive the various vectors of the present invention.

As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to bacterial enzymes, each of which cut double-stranded DNA at or near a specific nucleotide sequence.

A cell has been "transformed" or "transfected" by
15 exogenous or heterologous DNA when such DNA has been introduced inside the cell. The transforming DNA may or may not be integrated (covalently linked) into the genome of the cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With
20 respect to eukaryotic cells, a stably transformed cell is one in which the transforming DNA has become integrated into a chromosome so

that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA. A "clone" is a population of
5 cells derived from a single cell or common ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth in vitro for many generations.

A "heterologous" region of the DNA construct is an identifiable segment of DNA within a larger DNA molecule that is not
10 found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. In another example, heterologous DNA includes coding sequence in a construct
15 where portions of genes from two different sources have been brought together so as to produce a fusion protein product. Allelic variations or naturally-occurring mutational events do not give rise to a heterologous region of DNA as defined herein.

As used herein, the term "reporter gene" refers to a
20 coding sequence attached to heterologous promoter or enhancer

elements and whose product may be assayed easily and quantifiably when the construct is introduced into tissues or cells.

Transcriptional and translational control sequences are DNA regulatory sequences, such as promoters, enhancers, 5 polyadenylation signals, terminators, and the like, which provide for the expression of a coding sequence in a host cell.

The amino acids described herein are preferred to be in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the 10 desired functional property of immunoglobulin-binding is retained by the polypeptide. NH₂ refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide. In keeping with standard polypeptide nomenclature, *J Biol. Chem.*, 15 243:3552-59 (1969), abbreviations for amino acid residues are shown in the following Table of Correspondence:

Y	Tyr	tyrosine
G	Gly	glycine
F	Phe	Phenylalanine
20 M	Met	methionine
A	Ala	alanine

	S	Ser	serine
	I	Ile	isoleucine
	L	Leu	leucine
	T	Thr	threonine
5	V	Val	valine
	P	Pro	proline
	K	Lys	lysine
	H	His	histidine
	Q	Gln	glutamine
10	E	Glu	glutamic acid
	W	Trp	tryptophan
	R	Arg	arginine
	D	Asp	aspartic acid
	N	Asn	asparagine
15	C	Cys	cysteine

It should be noted that all amino-acid residue sequences are represented herein by formulae whose left and right orientation is in the conventional direction of amino-terminus to carboxy-terminus. Furthermore, it should be noted that a dash at the beginning or end of an amino acid residue sequence indicates a

peptide bond to a further sequence of one or more amino-acid residues. The above Table is presented to correlate the three-letter and one-letter notations which may appear alternately herein.

Thus, the present invention is directed to a fusion protein
5 comprising GFP so that the resulting fusion protein has a half life of
no more than about ten hours and as little as less than one hour. In
a preferred form, the GFP is EGFP. Preferably, the fusion protein
comprises EGFP fused to a PEST sequence-containing portion of a C-
terminus of murine ornithine decarboxylase (MODC). Representative
10 examples of PEST sequence-containing portion of a C-terminus of
murine ornithine decarboxylase include MODC₃₇₆₋₄₆₁, MODC₃₇₆₋₄₅₆,
MODC₄₂₂₋₄₆₁, P426A/P427A, P438A, E428A/E430A/E431A,
E444A, S440A, S445A, T436A, D433A/D434A and D448A. One
example of the GFP fusion protein of the present invention has the
15 sequence shown in SEQ ID No. 1.

The present invention is also directed to an isolated DNA
molecule encoding the fusion protein which comprises a fluorescent
protein selected from the group consisting of EGFP, ECFP and EYFP.
One example of the isolated DNA of the present invention has the
20 sequence shown in SEQ ID No: 2. The present invention is also
directed to a vector capable of expressing this isolated DNA molecule.

In one form, the vector contains a inducible promoter and is a tetracycline-regulated expression vector.

The present invention is also directed to a method of producing a stable cell line that expresses a fluorescent protein, e.g.,
5 GFP, comprising the step of transfecting cells with a vector disclosed herein.

In addition, the present invention is directed to a method of assaying activation or deactivation of promoters or other transcriptional or translational elements with a transient fluorescent
10 reporter protein, comprising the steps of transfecting cells with an expression vector comprising a fusion protein having a half life of no more than about ten hours, preferably less than four hours and most preferably less than one hour, wherein the fusion protein is under the influence of the promoter, transcriptional or translational
15 element, and detecting the presence, absence or amount of fluorescence in said cells. In this method, the amount of fluorescence present in the cell is a measure of the fluorescent protein that is being expressed. Detecting differences in fluorescence intensity between cells expressing the fluorescent protein under different
20 transcriptional or translational elements of interest is a rapid and straightforward procedure to measure effects of these transcriptional

or translational elements. Further, an additional step may be performed wherein transfected cells are treated with a compound of interest to determine the effect of the compound of interest on the transcriptional or translational elements. Detecting a change in
5 fluorescence upon treatment of the cells with the compound of interest is a rapid and straightforward procedure to measure the effects of the compounds on interest on the transcription or translation of the expressed fusion protein.

09931-081501
TOSTBO-2E2TE660
In addition, the present invention is directed to methods
10 of studying cell lineage comprising the steps of transfecting undifferentiated cells with a vector capable of expressing the destabilized fluorescent fusion protein of the present invention, growing the undifferentiated cells under conditions in which the undifferentiated cells become differentiated cells, and detecting an
15 absence or presence of fluorescence in the differentiated cells. Further, the present invention provides a method of using a fusion protein described herein in cell localization studies, comprising the steps of transfecting cells with an expression vector comprising a fluorescent fusion protein having a half life of no more than ten
20 hours and preferably less than four hours, wherein the fusion

protein is linked to a putative cell localization element, growing the cell and detecting a location of fluorescence in the cells.

EXAMPLE 1

5

Construction of DNA Expression Vectors

10 The cDNAs encoding EGFP and the C terminus of murine ODC (MODC) were amplified with pfu DNA polymerase (Stratagene, Inc., La Jolla, CA). EGFP was amplified with a pair of primers: 5' incorporated with a SacII recognition sequence and 3' with a Hind III sequence. The stop codon of EGFP was deleted from its C-terminus in order to make an open reading frame with the C terminus of murine ODC. The C terminus of murine ODC was also amplified with a pair of primers: 5' incorporated with a Hind III recognition sequence and 3' 15 with an EcoRI sequence. Two amplified PCR products were ligated at the Hind III site and the fusion was cloned into pTRE expression vector, Tc-regulated expression system (Gossen M., and Bujard H. (1992) *Proc. Natl. Acad. Sci.* 89: 5547-5551).

20 Using these methods, fusion proteins of EGFP-MODC were constructed. The EGFP-MODC₃₇₆₋₄₆₁ fusion protein included the complete C-terminus of murine ornithine decarboxylase. EGFP-

MODC₃₇₆₋₄₅₆ and EGFP-MODC₄₂₂₋₄₆₁ included only portions of the murine ornithine decarboxylase degradation domain, though both included the PEST sequence. Further, key amino acids of the PEST sequence in the fusion protein were then mutated to alanine using a
5 homology extension procedure (Rogers, et al., (1986) *Science* **234**, 364-368). The PEST mutants included P426A/P427A; P438A; E428A/E430A/E431A; E444A; S440A; S445A; T436A; D433A/D434A; and D448A.

10

EXAMPLE 2

Cell Transfection

15 The construct DNAs were purified and transfected into CHO K1-off cells for determination of protein degradation. CHO K1-off cells are CHO cells which were pre-transfected by a fusion protein of the tet-repressor and the herpes simplex virus VP16 gene (tTA). This pre-transfection allows expression of the gene coding for the
20 fusion protein on a pTRE vector (Gossen and Bujard, *ibid*), which in turn initiates transcription by binding to a modified CMV promoter

with tet-repressor binding elements. This binding can be blocked by tetracycline; hence, the expression can be controlled by tetracycline. The DNAs were introduced into these cells by CLONfectin (CLONTECH Laboratories, Inc., Palo Alto). After 24 hours, transfected cells were
5 subject to functional analyses.

EXAMPLE 3

Fluorescence Analysis

10 Cells were cultured on top of cover-slips to allow observation under a fluorescence microscope. After transfection, the cells were incubated at 37°C for 24 hours on the cover-slips and then fixed with 4% paraformaldehyde for 30 minutes. The cover-slips were mounted on a glass slide for fluorescence examination with a
15 Zeiss Axioskop Model 50 fluorescent microscope. To determine protein turnover, the cells were treated with cycloheximide at a final concentration of 100 µg/ml for variable times before paraformaldehyde fixation.

For FACS analysis, the transfected cells as well as
20 cycloheximide-treated cells were collected by EDTA treatment and the cell pellets were resuspended in 0.5 ml of PBS. The cell

suspensions were then analyzed for fluorescence intensity by FACS Calibur (Becton Dickson, Inc., San Jose, CA). EGFP was excited at 488nm, and emission was detected using a 510/20 bandpass filter.

5

EXAMPLE 4

Western Blot Analysis

For western blot analysis, transfected control cells as well
10 as cycloheximide-treated cells were collected in PBS and sonicated to
prepare cell lysates. Proteins were separated by SDS-PAGE. EGFP
and MODC fusion proteins were detected by a monoclonal antibody
against GFP after the proteins were transferred onto a membrane.
The detection was visualized with a chemiluminescent detection kit
15 (CLONTECH).

EXAMPLE 5

Determination of EGFP-MODC Protein Stability

20 The C terminus of murine ornithine decarboxylase, from
amino acids 376 to 461, has been shown to induce TbODC

degradation in mammalian cells. To demonstrate that the degradation domain could also induce EGFP degradation, murine ODC sequence was appended to the C-terminal end of EGFP to make a first fusion EGFP-MODC construct (Figure 1). The EGFP-MODC₃₇₆₋₄₆₁ fusion construct was expressed with the Tc-regulated expression vector (pTRE). Fluorescence intensity of the EGFP-MODC₃₇₆₋₄₆₁ fusion protein was examined under a fluorescence microscope after being transiently expressed in CHO K1-off cells. The fluorescence intensity of the EGFP-MODC₃₇₆₋₄₆₁ fusion protein was very low (Figure 1). Although it was believed that the lower fluorescence of the protein was due to rapid degradation, an EGFP fusion protein with such a low signal intensity would not be useful for most research applications.

If the rapid degradation indeed accounted for the lower fluorescence of the EGFP-MODC₃₇₆₋₄₆₁, the fluorescence intensity of the fusion protein could be increased by decreasing the rate of its degradation. The size of the C-terminal extension of murine ornithine decarboxylase can determine the rate of degradation. Deletion from either end of the degradation domain has yielded truncated proteins with a decreased rate of degradation, and removal

of the last five amino acids from murine ornithine decarboxylase dramatically decreases degradation of murine ornithine decarboxylase (Ghoda, L., et al. (1992) *Mol. Cell. Biol.* **12**, 2178-2185).

A TbODC fusion with a smaller extension starting at amino acid 422

5 has degraded more slowly than the longer extension starting at amino acid 376 (Li, X., and Coffino, P. (1993) *Mol. Cell. Biol.* **13**: 2377-2383). Therefore, two smaller extensions, one from amino acids 376

to 456 and the other from 422 to 461, also were appended to the C-terminus of EGFP to make EGFP-MODC₃₇₆₋₄₅₆ and EGFP-MODC₄₂₂₋

10 461 (Figure 1). Both of these fusion proteins contain the PEST sequence. After transfection, the fluorescence intensities of both

fusion proteins were examined with fluorescence microscopy.

Results indicated that both had a higher relative fluorescence intensity than EGFP-MODC₃₇₆₋₄₆₁, particularly the fusion protein

15 EGFP-MODC₄₂₂₋₄₆₁. As can be seen in Figure 1, the fluorescence of this latter fusion protein is similar to EGFP.

EXAMPLE 6

Further Characterization of EGFP-MODC₄₂₂₋₄₆₁ Protein Stability

Next, it was determined whether the C terminal extension
5 from amino acids 422 to 461 was able to induce EGFP degradation *in vivo*. To do this, the construct first was transiently transfected into CHO K1-off cells, and the half life of the fusion protein was determined by blocking protein synthesis with cycloheximide (CTX). At 24 hours post-transfection, the cells were treated with 100 µg/ml
10 cycloheximide for 0, 1, 2, 3, and 4 hours. The change in fluorescence intensity of the transfected cells was examined by fluorescence microscopy, and the results are shown in Figure 2. The fluorescence intensity of the fusion protein in the cells gradually decreased as cycloheximide treatment was extended, indicating that the EGFP-
15 MODC₄₂₂₋₄₆₁ fusion protein is unstable. After 4 hours of treatment with cycloheximide, less than half of the fluorescent intensity existed compared to the intensity of the fluorescence of the cells time zero. These results indicated that the half life of the fusion protein is less than 4 hours.

105130" 222FE660

The EGFP-MODC₄₂₂₋₄₆₁ was then compared to EGFP in the same assay. There was no significant change in the fluorescence intensity of EGFP in the EGFP-transfected cells four hours after protein synthesis stopped (Figure 2), indicating the half life of EGFP is longer than 4 hours. This is consistent with other reports on GFP. This result supports the conclusion that EGFP is a stable protein when expressed in mammalian cells and that the protein product of the EGFP-MODC₄₂₂₋₄₆₁ construct is unstable.

10 In order to determine more accurately the half lives of the EGFP-MODC fusion protein and EGFP, the change in fluorescence of these two proteins was quantified by flow cytometry. Transfected cells, after treatment with cycloheximide for 0, 1, 2 and 3 hours, were collected with EDTA, and 10,000 cells were subjected to FACS analysis. The results agreed with the fluorescence microscope observations; i.e., the fluorescence of the fusion protein gradually decreased as cycloheximide treatment was extended (Figure 3).

20 The graph in Figure 4 summarizes the FACS data and shows that approximately 50% of untreated cells maintained fluorescence after 2 hours of cycloheximide treatment, indicating that the half life of the fusion protein is about 2 hours. EGFP-transfected cells were subjected to similar analysis, and results

showed that the EGFP fluorescence did not change significantly during treatment with cycloheximide. Indeed, after 4 hours of cycloheximide treatment, EGFP cells still had more than 80% of fluorescence relative to untreated EGFP cells. In short, the half life of the EGFP fusion is suitably reduced and the half life of EGFP is significantly more than 4 hours.

Cycloheximide treatment cannot be prolonged for greater than 4 hours since it is toxic to cells, causing apoptosis. However, because the inducible expression system used in these studies is regulated by tetracycline (Gossen M., and Bujard H. (1992) *Proc. Natl. Acad. Sci.* 89: 5547-5551), EGFP synthesis can be stopped simply by adding tetracycline. To determine more precisely the half life of EGFP, the fluorescence intensities of EGFP transfected cells after 24 hours, in both the presence and absence of tetracycline, was monitored. EGFP first was allowed to be expressed for 24 hours after transfection. The transfected cells then were cultured in the presence or absence of tetracycline for another 24 hours and collected for analysis by flow cytometry. No difference in fluorescent intensity was detected between these two types of cells (-TC and +TC in the bottom panel of Figure 3), indicating that fluorescence did not

change in the 24 hours after EGFP protein synthesis shut-off. These results indicate that the half life of EGFP is more than 24 hours.

To examine if the half life of the EGFP and the EGFP fusion protein of the present invention correlated with the amount of fluorescence, degradation of the fusion protein was monitored by western blot analysis. Both the EGFP and EGFP-MODC₄₂₂₋₄₆₁ transfected cells that were used for flow cytometry in figure 3 were also used for Western blot analysis with a monoclonal antibody against GFP. As shown in figure 5, no detectable change in the level of EGFP protein was found among cells treated for 0-3 hours with cycloheximide, indicating that EGFP is stable during 3 hour cycloheximide treatment.

EGFP fusion proteins with the MODC modification of the present invention were also detected by a GFP monoclonal antibody. The size of the fusion protein was about 31 kDa. Unlike EGFP, however, the EGFP fusion protein was unstable. The level of the fusion protein fell dramatically by the end of the 3-hour cycloheximide treatment; in fact, less than one half of the control EGFP fusion protein was left at one hour, indicating that the half life of the fusion protein may be one hour or less. The difference in the measured half life using flow cytometry versus western blot analysis

likely is due to the fact that both premature (i.e. non-fluorescent) and mature GFP are detected by western analysis. However, the formation of EGFP chromophore is post-translational and proceeds with a half-time of about 25 minutes (Cormack, et al., (1996) *Gene* 173, 33-38). GFP in the context of this invention is important at its fluorescence level as a reporter, rather than at a protein level-- the half life of EGFP fluorescence is more important. The fluorescence half life of EGFP-MODC₄₂₂₋₄₆₁ *in vivo* is approximately 2 hours.

The amino acid sequence of the EGFP-MODC₄₂₂₋₄₆₁

10 protein is as follows:

MVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTGK
LPVPWPTLVTTLTYGVCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDG
NYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNVYIMADKQ
KNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSALSKD

15 PNEKRDHMLVLEFVTAAGITLGMDELYKKLSHGFPPEVEEQDDGTLPMSCAQ
ESGMDRHPAACASARINV (SEQ ID No:1); and the DNA sequence which encodes the EGFP-MODC₄₂₂₋₄₆₁ protein is as follows:

atgg tgagcaaggg cgaggagctg ttcaccgggg tgggtcccat cctggtcgag
ctggacggcg acgtaaacgg ccacaagttc agcgtgtccg gcgagggcga gggcgatgcc
20 acctacggca agctgaccct gaagttcatc tgcaccaccg gcaagctgcc cgtgccctgg
cccaccctcg tgaccaccct gacctacggc gtgcagtgtc tcagccgcta ccccgaccac

atgaagcagc acgacttctt caagtccgcc atgcccgaag gctacgtcca ggagcgcacc
atcttcttca aggacgacgg caactacaag acccgcgccg aggtgaagtt cgagggcgac
accctggtga accgcatcga gctgaagggc atcgacttca aggaggacgg caacatcctg
gggcacaagc tggagtacaa ctacaacagc cacaacgtct atatcatggc cgacaagcag
5 aagaacggca tcaaggtgaa cttcaagatc cgccacaaca tcgaggacgg cagcgtgcag
ctcgccgacc actaccagca gaacaccccc atcggcgacg gccccgtgct gctgcccgac
aaccactacc tgagcaccca gtccgccctg agcaaagacc ccaacgagaa gcgcgatcac
atggtcctgc tggagtctgt gaccgccgcc gggatcactc tcggcatgga cgagctgtac aag-
aagctt-agccatg gcttcccgcc ggaggtggag gagcaggatg atggcacgct gcccatgtct
10 tgtgcccagg agagcgggat ggaccgtcac cctgcagcct gtgcttctgc taggatcaat
gtgtagatgc (SEQ ID No:2).

EXAMPLE 7

Analysis of the PEST Sequence

The C terminus of mouse ornithine decarboxylase contains a PEST sequence from amino acids 423 to 449. There are three proline residues, four glutamic acid residues, two serine
20 residues, and one threonine residue (Figure 6). To evaluate the contribution of each amino acid in the PEST motif to the rate of

protein degradation, each Pro, Glu, Ser, and Thr residue in the PEST region of EGFP-MODC fusion protein was mutated to Ala. Degradation was monitored by the change of fluorescence. Each construct was transiently transfected into CHO K1-off cells. After treatment in the presence of cycloheximide for 0, 2, and 4 hours, the cells were collected for flow cytometry analysis. Data are shown in Figure 7.

Mutation of the proline residue at amino acid 438 stabilized the protein. After 4 hours of cycloheximide treatment, the percentage of fluorescent cells was still greater than 60%. Thus, the half life of the P438A mutant is more than double that of EGFP-MODC₄₂₂₋₄₆₁ suggesting that the existence of proline at 438 contributes to the instability of the fusion protein. Mutation of the prolines at amino acid positions 426 and 427 did not extend the half life. Instead, the half life of P426A/P427A is even shorter than that of EGFP-MODC₄₂₂₋₄₆₁, indicating that these proline residues may stabilize the protein. Similar results were obtained with mutations to the glutamine and serine in the fusion construct. The half lives of mutant E444A and S440A are longer than that of that EGFP-MODC₄₂₂₋₄₆₁, but E428A/E430A/E431A and S445A became more unstable, with only 20% or 29% of the cells retaining fluorescence after two hours of treatment.

105T80" 2E2TE660
10
The PEST sequence is often flanked by basic amino acids (Rogers, et al., (1986) *Science* 234, 364-368). To show the involvement of these flanking amino acid residues in protein instability, histidine 423, arginine 449 and histidine 450 were
5 mutated to alanine. Substitutions of alanine for the arginine and histidine at positions 449 and 450 dramatically increased protein stability, suggesting that these two amino acids are required for efficient protein degradation. Mutation of the His at amino acid 423 did not change protein stability.

EXAMPLE 8

Cell Line Expressing dEGFP With Tet Expression System

15 CHO K1-tet off cells were transfected with pTRE-EGFPMODC₄₂₂₋₄₆₁ and pTK-hygromycin. The transfection was performed with a CLONfectin kit (CLONTECH). The transfected cells were selected in the presence of 200 µg/ml hygromycin and resistant colonies were screened for fluorescence under a fluorescent
20 microscope. The individual single colonies of the fluorescent cells were transferred to new plates. As in the transiently transfected

cells, destabilized EGFP in the stably-transfected cells was regulated by tetracycline, and degradation was detectable by adding cycloheximide to block protein synthesis. The resulting stably-transfected cell line can be used for many analyses, including drug screening; in particular, it can be used to screen for drugs that block either the transcriptional induction of destabilized EGFP during the transition from the presence of tetracycline to the absence of tetracycline, or for drugs that inhibit protein degradation after the addition of cycloheximide.

EXAMPLE 9

Destabilized ECFP and EYFP

CFP (cyan) and YFP (yellow) are color variants of GFP. CFP and YFP contain 6 and 4 mutations, respectively. They are Tyr66Try, Phe66Leu, Ser65Thr, Asn145Ile, Met153Thr, and Val163Ala in CFP and Ser65Gly, Val68Leu, Ser72Ala, and Thr203Tyr. The enhanced CFP (ECFP) and YFP (EYFP) are encoded by genes with human-optimized codons. ECFP is excited at 433 nm and emits at 475 nm. EYFP is excited at 523 or 488 nm and emits at 527 nm.

Using the same strategy described above to create dEGFP, dECFP and dEYFP were generated by appending the C terminus of mouse ornithine decarboxylase to the C terminus of each of these two proteins (Figure 8). The half-lives of dECFP and dEYFP were
5 determined by transfecting pTRE-dECFP and pTRE-dEYFP into CHO/tTA cells, stopping protein synthesis by adding CHX, and subjecting to fluorescence microscopy. Both proteins were unstable in the presence of CHX. Two hours after treatment with CHX, half of fluorescence of these two proteins disappear indicating substantially
10 reduced half lives for these proteins as compared to wild type.

These two color variants of destabilized proteins can be used together for two-color detection. For example, the dECFP gene can be linked to the NFkB binding sequence and the dEYFP to the NFAT binding sequence. Introduction of these two reporter fusions
15 to a single cell allows detection simultaneous of two transcription factors or two signaling pathways simply by monitoring the relative fluorescence of the two colors.

EXAMPLE 10

Degradation Rate and Induction Fold of dEGFP

By mutating key amino acids of the PEST sequence of dEGFP, a number of mutants with different half lives were created. They were designated d1EGFP and d4EGFP, reflective of their half lives (Figure 9). The d1EGFP mutant has a half-life of less than 1 hour and the d4EGFP mutant has a half-life of approximately 4 hours (Figure 10). The EGFP-MODC₄₂₂₋₄₆₁ was designated d2EGFP because its half-life is about 2 hours.

The outstanding character of dEGFP proteins, when compared to EGFP, is that they exhibit a rapid turnover resulting in less accumulation of the destabilized proteins even at basal-level activity. Therefore, d2EGFP is a much more sensitive transcription reporter than EGFP, as has been demonstrated with the system of TNF-mediated NFkB activation. Because d1EGFP has a shorter half-life than d2EGFP, d1EGFP is more sensitive than d2EGFP when used as a transcription reporter. To illustrate this, d1EGFP and d2EGFP genes were linked to cAMP regulated element (CRE) and their responses to forskolin for each fusion construct tested. Four-fold induction was obtained when d2EGFP was used as the reporter,

however more than 12-fold induction was achieved when d1EGFP used (Figure 11). These results indicate that a more rapid turnover of the GFP molecule achieves higher induction and enhanced sensitivity.

5 Any patents or publications mentioned in this specification are indicative of the levels of those skilled in the art to which the invention pertains. These patents and publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to
10 be incorporated by reference.

One skilled in the art will readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The present examples along with the methods, procedures, treatments, molecules, and specific compounds described herein are
15 presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention as defined
20 by the scope of the claims.